

*Sperm-depleted males influence the reproductive behaviour of conspecifics*

**Philippe Louâpre, Stéphanie Llopis,  
Véronique Martel & Joan van Baaren**

**Naturwissenschaften**  
The Science of Nature

ISSN 0028-1042  
Volume 101  
Number 11

Naturwissenschaften (2014)  
101:875-882  
DOI 10.1007/s00114-014-1227-6



**Your article is protected by copyright and all rights are held exclusively by Springer-Verlag Berlin Heidelberg. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at [link.springer.com](http://link.springer.com)".**

# Sperm-depleted males influence the reproductive behaviour of conspecifics

Philippe Louâpre · Stéphanie Llopis · Véronique Martel · Joan van Baaren

Received: 10 April 2014 / Revised: 9 August 2014 / Accepted: 11 August 2014 / Published online: 3 September 2014  
© Springer-Verlag Berlin Heidelberg 2014

**Abstract** In many insect species, sperm-depleted males (SDMs, i.e. males that have exhausted their sperm after a given number of matings) remain sexually active and continue to mate females. Here, we investigated the behavioural modifications that occur in both sexes of the parasitoid *Asobara tabida* Nees (Hymenoptera: Braconidae), after matings by fertile males and sperm-depleted males. We show that (i) virgin females, mated females and females mated to a SDM exhibited different behaviours and that (ii) males responded differently to females depending on whether the females had previously mated with an SDM or not. Our findings demonstrate that SDM influenced the reproductive behaviour of both males and females, especially with regard to male responsiveness and female attractiveness. These findings are discussed in the context of adaptive behaviour and fitness maximization in both males and females.

**Keywords** Sperm-depleted male · Hymenopteran parasitoids · Mate choice · *Asobara tabida* · Conspecific behaviour

## Introduction

Sperm is not an infinite and free-cost resource to produce. In polygynous species, mating several females successively in a short period of time may result in sperm depletion: The sperm supply can be drained after a given number of females mated (Wedell et al. 2002). Sperm depletion can be temporary (in synspermatogenic species) or definitive (in prospermatogenic species), depending on the ability of males to produce new sperm during their adult life (Damiens and Boivin 2006; Boivin 2013). Sperm-depleted males (SDMs) have been found in many taxa, such as mammals (Preston et al. 2001), birds (Pizzari et al. 2003), lizards (Olsson et al. 2004), crustaceans (Rubolini et al. 2007) and fish (Smith et al. 2009). In insects, sperm depletion occurs in many species, and SDM individuals remain sexually active as they are still able to attract receptive females and to mate them (Simmonds 1953; Laing and Caltagirone 1969; Gordh and DeBach 1976; Van Den Assem et al. 1979 Nadel and Luck 1985; Ramadan et al. 1991; Ode et al. 1995; Damiens and Boivin 2005; Rönn et al. 2008; Steiner et al. 2008). It was suggested that continuing to mate is an adaptive strategy of SDMs to decrease the fitness of fertile males (Damiens and Boivin 2006). The authors indeed demonstrated that *Trichogramma evanescens* Westwood (Hymenoptera: Trichogrammatidae) females have a decreased ability to store sperm after being mated by SDMs (Damiens and Boivin 2006). To continue to mate is thus viewed as a SDMs' strategy to increase their relative fitness comparing to this of other males. On the other hand, such behaviour may persist over generations because of the lack of selective pressure counter-selecting it (SDMs can obviously not produce

Communicated by: Sven Thatje

Véronique Martel and Joan van Baaren have contributed equally to the manuscript and are thus co-last authors.

P. Louâpre (✉)  
Biodiversity Research Centre, Earth and Life Institute,  
Université Catholique de Louvain, 4-5, Place Croix du Sud B-1348  
Louvain-la-Neuve, Louvain-la-Neuve, Belgium  
e-mail: philippe.louapre@mail-univscience.com

P. Louâpre · S. Llopis · J. van Baaren  
UMR CNRS 6553 Ecologie, Biodiversité, Evolution,  
Université de Rennes 1, Rennes, France

S. Llopis  
e-mail: stephanie.llopis@univ-rennes1.fr

J. van Baaren  
e-mail: joan.van-baaren@univ-rennes1.fr

V. Martel  
Natural Resources Canada, Canadian Forest Service,  
Laurentian Forestry Centre, Alberta, Canada  
e-mail: veronique.martel@nrcan-rncan.gc.ca

progeny anymore). Whatever the adaptive significance of the SDM mating behaviour, its effect on the behaviours of conspecifics (both males and females) remains largely unknown.

Females' behaviours may be modified after copulation; after mating, females usually become refractory to subsequent copulations for a given time, even without sperm transfer (Gillott 2003). This refractoriness to mating is caused by chemicals contained in the seminal fluids transferred during mating or by the male's behaviour (Alcock 1994; Gillott 2003; Wedell 2005; Yamane 2013). The transfer of chemicals to the mated female is therefore viewed as a male strategy to suppress the subsequent reproductive activity of females, thus decreasing the likelihood of sperm competition (Ejima et al. 2007). For example, Radhakrishnan et al. (2009) showed that the transfer of seminal fluid in *Bactrocera tryoni* Froggatt (Diptera: Tephritidae) during mating induces female sexual inhibition. In this case, the behaviour of the mated females is modified by SDM resulting in fewer opportunities for other males to access them. In many species, females mated to a SDM become less receptive to mating than virgin females, suggesting that these females no longer invest in pheromone biosynthesis after mating (Steiner et al. 2008). We however note that in at least another species, females mated to a SDM remain sexually active (Damiens and Boivin 2006).

Male mating may also modify the behaviour of conspecific males. Males are known to be choosy just like females (Wedell et al. 2002), and the females' mating status influences male choice. For example, the male preferences for virgin females and/or for females that they have never previously encountered have been observed in many insect species (Bonduriansky 2001; Wedell et al. 2002; Ferveur 2005; Bateman and Fleming 2006; Everaerts et al. 2010). Sexual pheromones transferred by males during mating can mark females as mated, making them less attractive to other males (Findlay et al. 2008). In *Drosophila*, mating induces changes in the female cuticular profile, which reduces the time spent in copulation by subsequent males (Friberg 2006; Everaerts et al. 2010). Previous mating can therefore influence the choice of other males, which generally prefer to mate with virgin females.

Because all the behavioural modifications detailed above do not require sperm transfer, such modifications are likely to occur after mating by a SDM. The aim of our study was thus to investigate the behavioural consequences of SDMs' mating behaviour in both sexes. In this system, conflict between the sexes arises, as females mated to a SDM are still unfertilized and so would gain by remaining attractive to other males in order to ensure egg fertilization. On the other hand, SDMs would gain by decreasing the attractiveness of the females they mated in order to reduce other males' opportunity to access them. Given that, we hypothesize that (i) females mated to a SDM should be as receptive as virgin females in order to get access to further insemination and (ii) males

should be more attracted to virgin females than females mated to a SDM and mated females.

To test these hypotheses, we used the parasitoid *Asobara tabida* Nees (Hymenoptera: Braconidae), a haplodiploid species in which sons result from unfertilized eggs while daughters result from fertilized eggs, allowing us to determine if the females had been successfully inseminated from the resulting sex ratio obtained—the presence of at least one daughter in the progeny is a proof of insemination. *A. tabida* males exhibit a stereotyped courtship behaviour consisting of wing fanning, antennation, head movements, and leg tapping (Matthews 1975; Carton et al. 1986; Dufour et al. 2012). No postcopulatory behaviour is reported for this species (Stökl et al. 2014). *A. tabida* males are generally sperm-depleted after a large number of matings (about 13 matings) but continue to mate after sperm depletion (Louâpre, Dufour, personal observation). Chemical communication is of great importance for mating as *A. tabida* males respond to chemicals deposited by females and increase their residence time in a patch previously exposed to females (Dufour et al. 2012). More recently, Stökl et al. (2014) highlighted the remarkable diversity of the sex pheromones of males and females in *A. tabida*. They showed that mating behaviour in this species is regulated by a complex combination of chemicals rather than by a single molecule.

## Materials and methods

### Insect rearing and general methods

The *A. tabida* strain used in this study was collected in Chasse-Sur-Rhône (France, 45°34'N, 4°48'E) in September 2008 and reared in the laboratory on second-instar *Drosophila subobscura* hosts in a controlled climate room (20 °C, 60 % relative humidity, and 16:8 h light/dark). A few days before emergence, parasitized pupae from the culture were isolated in gelatin capsules (0.68 mL) to obtain virgin parasitoids of known age. Freshly emerged males and females were kept separately in plastic tubes ( $h=10$  cm,  $\varnothing=2.5$  mm) and fed with diluted honey solution.

### Choice tests

Experiments were performed in the controlled climate room used for the rearing. Males were 1–2 days old, and females were an equal mix of 1–4 days old (the two females used in each if the choice test were of comparable ages in order to avoid potential confounding effect of age). A light table was used to maintain the individuals on the bottom of a petri dish ( $\varnothing=90$  mm) by phototaxis. All experiments were videotaped for a period of 15 min and analyzed using the computer software package The Observer v3.0 (Noldus, Wageningen,

The Netherlands). During each choice test, we noted (i) the male behaviour (courtship, contact with the female, mating), (ii) the identity of the female towards which the male directed its behaviour, and (iii) the behaviour of the female towards the male (i.e. approaches and escapes, acceptance or rejection). The identity of the female towards which the first courtship was directed was also noted as a proxy of the male preference before any reaction from the female. Mating is indeed the result of both active and apparent choices of the two sexes, and the first courtship give supplementary information compared to the mating with a given female (Martel and Boivin 2011).

#### Experiment 1: female mated to a SDM vs virgin female

In this experiment, a virgin male was placed in the presence of both a female mated to a SDM and a virgin female in order to find out whether the male would perceive them as different resources. To obtain a female mated to a SDM, a single male mated a sequence of 30 virgin females over a period of 6 h (from rank 1 to rank 30 females). Although *A. tabida* males become sperm-depleted after about 13 matings (Louâpre, Dufour, personal communication), we used females from rank 24 to rank 29 and we isolated the rank 23 and rank 30 females in a plastic tube containing a substrate of agar-nipagine and 20-s instar *D. subobscura* larvae. The sex ratio produced was noted, and only data from true SDMs and females mated to an actual SDM—whose offspring had a sex ratio of 1 (i.e. male-only progeny)—were used for the analysis. Under the rearing conditions, inseminated females indeed laid at least several females in a dozen of hosts, their sex ratio being normally female-biased, so a male-only progeny obtained in 20 hosts was used as an indicator for an absence of sperm transfer to the females. One female mated to a SDM (only females mated to different SDMs were used for each of the repetitions) and virgin female were then placed in the same clean petri dish. A virgin male was introduced at the beginning of the experiment and placed equidistant from the two females. Thirteen dual choice tests were performed.

#### Experiment 2: female mated to the same SDM vs virgin female

In this experiment, each SDM was placed in the presence of a female he had previously mated without sperm transfer (randomly selected from rank 24 to rank 29 in the sequence of copulations) and a virgin female. Both the female mated to the SDM and the virgin female were placed in the same clean petri dish. The SDM was introduced at the beginning of the experiment and placed equidistant from the two females. Eleven dual choice tests were performed.

#### Experiment 3: female mated to the same fertile male vs female mated to another fertile male

In this experiment, a fertile male was placed in the presence of a female he had previously mated and a female that had been mated by another male to determine whether males recognize a female they have previously mated. Ten minutes before the choice test, two virgin males each mated a virgin female in separate plastic tubes ( $h=10$  cm,  $\varnothing=2.5$  mm). The two mated females were then placed in the same clean petri dish. One of the two males used was introduced at the beginning of the experiment and placed equidistant from the two females. Twenty-two dual choice tests were performed.


#### Statistical analyses

Statistical analyses were performed using R 2.15.0 software at the critical level  $\alpha=0.05$ . The male choices were compared to a 50:50 probability (i.e. no preferential choice) using an exact binomial test. The mean numbers of the types of behaviour exhibited by the males and females were analyzed using Wilcoxon tests. Because both virgin females and females mated to a SDM were used in the first two experiments, the occurrence of approaches and escapes during these two experiments were analyzed by generalized linear model (GLM)-Poisson procedure, using the mating status of the females (i.e. virgin, pseudo-virgin or mated) and of the males (fertile or sperm-depleted) as factors. Multiple comparisons were then performed using the `glt` function of R (`multcomp` package).

## Results

#### Experiment 1: female mated to a SDM vs virgin female

The two types of females exhibited the same number of approaches and escapes towards the virgin male (respectively,  $W=82$ ,  $P=0.915$  and  $W=78$ ,  $P=0.753$ ) (Fig. 1b). Virgin males did not direct their first courtship or their first mating towards one female type rather than the other (binomial test, respectively,  $P=0.581$  and  $P=0.179$ ) (Fig. 1a). During the course of the experiment, the males exhibited on average the same number of courtships and contacts with females mated to a SDM and virgin females (respectively,  $W=68$ ,  $P=0.401$  and  $W=64$ ,  $P=0.288$ ) (Fig. 1c). They tended to mate the originally virgin females more frequently than the SDM-mated females during the course of the experiment, but this difference was not significant probably due to the weak number of repetitions ( $W=52.5$ ,  $P=0.055$ ) (Fig. 1c).



**Fig. 1** Behaviours of virgin males (**a**, **b**), virgin females and females mated to a SDM (**c**) during the first experiment. (**a**) Proportion ( $\pm 95\%$  confidence interval) of virgin males exhibiting their first courtship towards (*top*,  $n=13$ ) and/or mated first with (*bottom*,  $n=9$ ) the proposed

females. (**b**) Mean number ( $\pm$ SE) of behaviour exhibited by virgin males towards different types of females. (**c**) Mean number ( $\pm$ SE) of behaviour exhibited by females towards virgin males (*ns* not significant)

#### Experiment 2: female mated to the same SDM vs virgin female


Virgin females exhibited more approaches towards the SDM than the females previously mated by him ( $W=26.5$ ,  $P=0.026$ ), while the number of escapes from the SDM did not differ between both groups of females ( $W=46$ ,  $P=0.352$ ) (Fig. 2b). The first courtship was not significantly exhibited by SDMs towards one type of females over the other (binomial test,  $P=0.227$ ) (Fig. 2a). On average, SDMs directed a similar number of courtships towards females mated by them and virgin females ( $W=35$ ,  $P=0.401$ ) (Fig. 2c). However, they exhibited more contacts with virgin females than with females they previously mated ( $W=29$ ,  $P=0.037$ ). They mated the virgin females before the females they previously mated (binomial test,  $P=0.012$ ; Fig. 2a) and mated the originally virgin females more frequently than the SDM-mated females during the course of the experiment, even though she becomes mated after the first mating ( $W=13.5$ ,  $P<0.001$ ; Fig. 2c).

#### Experiment 3: female previously mated by the same fertile male vs female previously mated by another fertile male

The two types of females exhibited the same number of approaches and escapes towards the fertile male ( $W=299$ ,  $P=0.104$  and  $W=199$ ,  $P=0.313$ , respectively) (Fig. 3b). Although fertile males directed their first courtship towards females previously mated by another male more often than towards the ones they had mated themselves (binomial test,  $P=0.004$ ) (Fig. 3a), they exhibited overall the same number of courtships and contacts towards both females ( $W=184.5$ ,  $P=0.171$  and  $W=211$ ,  $P=0.458$ , respectively) (Fig. 3c). No mating was observed during this experiment.

#### Female behaviour across the experiments

The behaviour of virgin females and females mated to a SDM was compared between experiments 1 and 2. The number of approaches by females differed depending on their mating



**Fig. 2** Behaviours of SDMs (a, b), virgin females and females mated to the same SDM (c) during the second experiment. (a) Proportion ( $\pm 95\%$  confidence interval) of SDMs exhibiting their first courtship towards (top,  $n=11$ ) and/or mated first with (bottom,  $n=11$ ) the proposed females. (b)

Mean number ( $\pm$ SE) of behaviour exhibited by SDMs towards different types of females. (c) Mean number ( $\pm$ SE) of behaviour exhibited by females towards the SDM (ns not significant;  $*P<0.05$ )


status (GLM-Poisson,  $F_{1,80}=4.007$ ,  $P=0.045$ ) but not on the male mating status (GLM-Poisson,  $F_{1,79}=0.442$ ,  $P=0.51$ ) (Fig. 1a, b): Females mated to a SDM exhibited more approaches than virgin females, regardless of the mating status of the males ( $z=-1.997$ ,  $P=0.046$ ). The number of escapes differed between females but differently depending on the mating status of the male (GLM-Poisson, female mating status: male mating status,  $F_{3,78}=2.992$ ,  $P=0.029$ ) (Fig. 1a, b): Females mated to a SDM exhibited the same number of escapes from virgin males and SDMs ( $z=-0.205$ ,  $P=0.997$ ), while virgin females exhibited more escapes from SDMs than virgin males ( $z=2.624$ ,  $P=0.043$ ).

## Discussion

We showed that (i) virgin females, mated females, and females mated to a SDM exhibited different behaviours and that (ii)

the mating status of the females influences the behaviours of males. Different behaviours were observed according to the reproductive status of the two sexes, although the majority of the females were of the same mating status at the end of the experiment. As pointed out by Martel and Boivin (2011), mating depends not only on the choice of the individual (true choice) but also on the response of the partner (apparent choice). Accordingly, we showed that the mating pattern in this species can be influenced by the SDM's behaviour, which, in turn, impacts both males' and females' choice. Our results thus demonstrate that SDMs influenced the male responsiveness and female attractiveness.

Sperm-depleted males influence the subsequent behaviour of both the females they have mated and virgin females. Females mated to a SDM were more attracted to males than virgin females whatever the reproductive status of the males. Moreover, virgin females exhibited more escapes from SDMs than from virgin males. Taken together, these findings show that (i) females mated to a SDM were more reproductively



**Fig. 3** Behaviours of fertile males (**a, b**), females mated to the same male or to another male (**c**) during the second experiment. (**a**) Proportion ( $\pm 95\%$  confidence interval) of fertile males exhibiting their first courtship towards ( $n=22$ ) the proposed females. (**b**) Mean number ( $\pm SE$ ) of

behaviour exhibited by the SDM towards different types of females. (**c**) Mean number ( $\pm SE$ ) of behaviour exhibited by females towards the SDM. (*ns* not significant;  $**P < 0.01$ )

active than virgin females and that (ii) SDMs are not as attractive for females as virgin males. Females mated to a SDM should be at least as active as virgin females in order to compensate for the unsuccessful insemination (Arnqvist and Nilsson 2000; Hosken et al. 2001; Chapman et al. 2003; Tregenza et al. 2006). Our results are consistent with this and demonstrate that even if SDMs transferred seminal fluid to the females, this did not inhibit female sexual receptivity as it was demonstrated for other species (Leopold 1976; Avila et al. 2011). Female *A. tabida* mated to a SDM are thus receptive to mating, as has been shown for female *T. evanescens* mated to a SDM (Damiens and Boivin 2006). Moreover, females mated to a SDM seemed to exhibit adaptive behaviour, because they reduced the number of approaches they made towards SDMs; whatever the mechanism behind this postmating effect, females mated to a SDM behave as if they can ‘learn’ and consequently modify their behaviour. This is comparable to other types of behaviour exhibited by females in order to

ensure that their eggs are fertilized (reviewed by Rhainds 2010).

The behaviour of males is also influenced by the reproductive status of the females; virgin females, females mated by a SMD and mated females do not elicit the same behavioural response from males according to the mating status of the males. Our results did not show any significant effect of the reproductive status of the females on the behaviour of virgin males (experiment 1). We however noted a tendency of virgin males to exhibit their first courtship and mating towards virgin females, and the small sample size could have led to non-significant results. SDMs, however, did exhibit their first courtship and mating towards virgin females over females they previously mated. They also exhibited more contacts towards virgin females over females they mated (experiment 2). These results demonstrate that females mated to a SDM are not perceived as virgin females by the males. The preferential choice of SDMs for virgin females rather than females mated

to a SDM—even though females mated to a SDM continue to be receptive to mating—suggests a decrease in the attractiveness of females mated to a SDM.

Many cues can be used by males to discriminate between virgin females and females mated to a SDM, and chemicals must play a major role (Thomas 2011). In bumblebees and parasitic wasps, sexual pheromones are indeed known to carry information about mating status (Lloyd 1981; Ode et al. 1995). In *Drosophila*, the cuticular profiles of females change after mating (Steiger et al. 2008). Moreover, the mating behaviour of *A. tabida*—like many other insect species—is based on a complex and dynamic infochemical emission and on the perceptions of both males and females (Stöckl et al. 2014). *A. tabida* males may thus perceive and respond to chemical changes in the mated females (with or without sperm being stored), which enable them to discern the female's reproductive status.

Males could be more attracted to virgin females than to females mated to a SDM because they never encountered the virgin females they faced with (the so-called 'Coolidge effect') (Wilson et al. 1963; Dewsbury 1981). Based on individual marking recognition, Steiner et al. (2008) have shown that males of the burying beetle *Nicrophorus vespilloides* Herbst (Coleoptera: Silphidae) use female cuticular patterns to discriminate between novel females and those they have previously inseminated. In our study, such a Coolidge effect may be responsible for the choice of males faced by two mated females (experiment 3) and the choice of a SDM facing virgin females and females mated to a SDM (experiment 1) but not for that of virgin males in the first experiment. Indeed, the virgin males used in our study had never encountered any females. The change in the behaviour of the males and females may therefore involve different mechanisms and does not depend on sperm transfer.

To conclude, our study showed that the mating process in *A. tabida* depends on the behaviour of both sexes and is not a simple and unidirectional choice of one sex towards the other. *A. tabida* males thus actively choose their mates on the basis of mating status, but females can accept or reject the interested males. SDMs and females mated to a SDM behave in different ways towards fertile males and virgin or mated females, respectively. As other authors have suggested, the modification of male and female behaviour as a result of mating could be a way of improving the fitness of SDMs (Damiens and Boivin 2006) by decreasing the likelihood of subsequent mating by other males through sperm competition avoidance. Many hypotheses have been advanced to explain the female mating failure in nature, including a lack of opportunity for mating (Rhains 2010). The mating behaviour of SDMs may contribute to this phenomenon by making females mated to a SDM look like successfully mated females to other males.

**Acknowledgments** We are grateful to Elise Blondelot and Myriam Labrecque for their technical help and Monica Gosh for English revision. This research was supported by a grant from the Ministère de l'Enseignement Supérieur et de la Recherche. All experiments conducted in this study comply with the current animal experimentation regulations in France.

## References

- Alcock J (1994) Postinsemination associations between males and females in insects: the mate-guarding hypothesis. *Ann Rev Entomol* 39: 1–21. doi:10.1146/annurev.en.39.010194.000245
- Arnqvist G, Nilsson T (2000) The evolution of polyandry: multiple mating and female fitness in insects. *Anim Behav* 60:145–164. doi:10.1006/anbe.2000.1446
- Avila FW, Sirot LK, LaFlamme BA et al (2011) Insect seminal fluid proteins: identification and function. *Annu Rev Entomol* 56:21–40
- Bateman PW, Fleming PA (2006) Males are selective too: mating, but not courtship, with sequential females influences choosiness in male field crickets (*Gryllus bimaculatus*). *Behav Ecol Sociobiol* 59: 577–581. doi:10.1007/s00265-005-0083-y
- Boivin G (2013) Sperm as a limiting factor in mating success in Hymenoptera parasitoids. *Entomol Exp App* 146:149–155. doi:10.1111/j.1570-7458.2012.01291.x
- Bonduriansky R (2001) The evolution of male mate choice in insects: a synthesis of ideas and evidence. *Biol Rev* 76:305–339. doi:10.1017/S1464793101005693
- Carton Y, Bouletreau M, Van Alphen JJM, Van Lenteren JC (1986) The *Drosophila* parasitic wasps. *Genet. Biol. Drosoph.*, M Ashburne. Academic Press, New York, pp 347–394
- Chapman T, Arnqvist G, Bangham J, Rowe L (2003) Sexual conflict. *Trends Ecol Evol* 18:41–47
- Damiens D, Boivin G (2005) Male reproductive strategy in *Trichogramma evanescens*: sperm production and allocation to females. *Physiol Entomol* 30:241–247. doi:10.1111/j.1365-3032.2005.00453.x
- Damiens D, Boivin G (2006) Why do sperm-depleted parasitoid males continue to mate? *Behav Ecol* 17:138–143. doi:10.1093/beheco/arj009
- Dewsbury DA (1981) Effects of novelty of copulatory behavior: the Coolidge effect and related phenomena. *Psych Bull* 89:464–482. doi:10.1037/0033-2909.89.3.464
- Dufour CM-S, Louâpre P, van Baaren J, Martel V (2012) When parasitoid males make decisions: information used when foraging for females. *PLoS One* 7:e46706. doi:10.1371/journal.pone.0046706
- Ejima A, Smith BPC, Lucas C et al (2007) Generalization of courtship learning in *Drosophila* is mediated by cis-vaccenyl acetate. *Curr Biol* 17:599–605. doi:10.1016/j.cub.2007.01.053
- Everaerts C, Farine J-P, Cobb M, Ferveur J-F (2010) *Drosophila* cuticular hydrocarbons revisited: mating status alters cuticular profiles. *PLoS One* 5:e9607. doi:10.1371/journal.pone.0009607
- Ferveur J-F (2005) Cuticular hydrocarbons: their evolution and roles in *Drosophila* pheromonal communication. *Behav Genet* 35:279–295. doi:10.1007/s10519-005-3220-5
- Findlay GD, Yi X, MacCoss MJ, Swanson WJ (2008) Proteomics reveals novel *Drosophila* seminal fluid proteins transferred at mating. *PLoS Biol* 6:e178. doi:10.1371/journal.pbio.0060178
- Friberg U (2006) Male perception of female mating status: its effect on copulation duration, sperm defence and female fitness. *Anim Behav* 72:1259–1268. doi:10.1016/j.anbehav.2006.03.021
- Gillott C (2003) Male accessory gland secretions: modulators of female reproductive physiology and behavior. *Ann Rev Entomol* 48:163–184. doi:10.1146/annurev.ento.48.091801.112657

- Gordh G, DeBach P (1976) Male inseminative potential in *Aphytis lingnanensis* (Hymenoptera: Aphelinidae). *Can Entomol* 108:583–589
- Hosken DJ, Garner TWJ, Ward PI (2001) Sexual conflict selects for male and female reproductive characters. *Curr Biol* 11:489–493
- Laing D, Caltagirone L (1969) Biology of *Habrobracon lineatellae* (Hymenoptera: Braconidae). *Can Entomol* 101:135–142
- Leopold R (1976) The role of male accessory glands in insect reproduction. *Annu Rev Entomol* 21:199–221
- Lloyd JE (1981) Sexual selection: individuality, identification, and recognition in a bumblebee and other insects. *The Florida Entom* 64:89. doi:10.2307/3494603
- Martel V, Boivin G (2011) Do choice tests really test choice? *J Insect Behav* 24:329–336. doi:10.1007/s10905-011-9257-9
- Matthews RW (1975) Courtship in parasitic wasps. *Evol. Strateg. Parasit. insects mites*. Springer, pp 66–86
- Nadel H, Luck R (1985) Span of female emergence and male sperm depletion in the female-biased, quasi-gregarious parasitoid, *Pachycrepoideus vindemiae* (Hymenoptera: Pteromalidae). *Ann Entomol Soc Am* 78:410–414
- Ode PJ, Antolin MF, Strand MR (1995) Brood-mate avoidance in the parasitic wasp *Bracon hebetor* Say. *Anim Behav* 49:1239–1248. doi:10.1006/anbe.1995.0156
- Olsson M, Madsen T, Ujvari B, Wapstra E (2004) Fecundity and MHC affect ejaculation tactics and paternity bias in sand lizards. *Evol* 58:906–909
- Pizzari T, Cornwallis CK, Løvlie H et al (2003) Sophisticated sperm allocation in male fowl. *Nature* 426:70–74. doi:10.1038/nature02004
- Preston BT, Stevenson IR, Pemberton JM, Wilson K (2001) Dominant rams lose out by sperm depletion. *Nature* 409:681–682. doi:10.1038/35055617
- Radhakrishnan P, Pérez-Staples D, Weldon CW, Taylor PW (2009) Multiple mating and sperm depletion in male Queensland fruit flies: effects on female remating behaviour. *Anim Behav* 78:839–846. doi:10.1016/j.anbehav.2009.07.002
- Ramadan MM, Wong TT, Wong MA (1991) Influence of parasitoid size and age on male mating success of *Opiinae* (Hymenoptera: Braconidae), larval parasitoids of fruit flies (Diptera: Tephritidae). *Biol Control* 1:248–255. doi:10.1016/1049-9644(91)90074-A
- Rhains M (2010) Female mating failures in insects. *Entom Exp App* 136:211–226. doi:10.1111/j.1570-7458.2010.01032.x
- Rönn JL, Katvala M, Arnqvist G (2008) Interspecific variation in ejaculate allocation and associated effects on female fitness in seed beetles. *J Evol Biol* 21:461–470. doi:10.1111/j.1420-9101.2007.01493.x
- Rubolini D, Galeotti P, Pupin F et al (2007) Repeated matings and sperm depletion in the freshwater crayfish *Austropotamobius italicus*. *Freshw Biol* 52:1898–1906. doi:10.1111/j.1365-2427.2007.01814.x
- Simmonds F (1953) Observations on the biology and mass-breeding of *Spalangia drosophilae* Ashm. (Hymenoptera, Spalangiidae), a parasite of the fruit-fly, *Oscinella frit* (L.). *Bull Entomol Res* 44:773–778
- Smith C, Pateman-Jones C, Zięba G et al (2009) Sperm depletion as a consequence of increased sperm competition risk in the European bitterling, *Rhodeus amarus*. *Anim Behav* 77:1227–1233
- Steiger S, Franz R, Eggert A-K, Müller JK (2008) The Coolidge effect, individual recognition and selection for distinctive cuticular signatures in a burying beetle. *Proc R Soc B* 275:1831–1838. doi:10.1098/rspb.2008.0375
- Steiner S, Henrich N, Ruther J (2008) Mating with sperm-depleted males does not increase female mating frequency in the parasitoid *Lariophagus distinguendus*. *Entom Exp App* 126:131–137. doi:10.1111/j.1570-7458.2007.00641.x
- Stökl J, Dandekar A-T, Ruther J (2014) High chemical diversity in a wasp pheromone: a blend of methyl 6-methylsalicylate, fatty alcohol acetates and cuticular hydrocarbons releases courtship behavior in the drosophila parasitoid *Asobara tabida*. *J Chem Ecol* 40:159–168. doi:10.1007/s10886-014-0378-4
- Thomas ML (2011) Detection of female mating status using chemical signals and cues. *Biol Rev* 86:1–13. doi:10.1111/j.1469-185X.2010.00130.x
- Tregenza T, Wedell N, Chapman T (2006) Introduction. Sexual conflict: a new paradigm? *Phil Trans R Soc B* 361:229–234. doi:10.1098/rstb.2005.1796
- Van Den Assem J, Gijswijt MJ, Nübel BK (1979) Observations on courtship and mating strategies in a few species of parasitic wasps (Chalcidoidea). *Netherlands J Zool* 30:208–227
- Wedell N (2005) Female receptivity in butterflies and moths. *J Exp Biol* 208:3433–3440. doi:10.1242/jeb.01774
- Wedell N, Gage MJG, Parker GA (2002) Sperm competition, male prudence and sperm-limited females. *Trends Ecol Evol* 17:313–320. doi:10.1016/S0169-5347(02)02533-8
- Wilson JR, Kuehn RE, Beach FA (1963) Modification in the sexual behavior of male rats produced by changing the stimulus female. *J Comp Physiol Psychol* 56:636–644
- Yamane T (2013) Reduction of female mating receptivity after injection of male-derived extracts in *Trigonotylus caelestialium* (Heteroptera: Miridae). *Ann Entomol Soc Am* 106:258–264. doi:10.1603/AN12079